

# Mylan On Location™

## Anaphylaxis and Administering EpiPen® (epinephrine injection) Auto-Injector Quiz

Name: \_\_\_\_\_

### 1. What is anaphylaxis?

- a. Any allergic reaction
- b. Inflammation of the blood vessels
- c. A life-threatening allergic reaction
- d. None of the above

### 2. Anaphylaxis signs and symptoms may:

- a. Vary from one event to the next
- b. Occur quickly (within 5 minutes) or progress over hours
- c. Both a and b
- d. None of the above

### 3. Anaphylaxis signs and symptoms may include:

- a. Tongue and/or lip swelling
- b. Hives and/or swelling
- c. Vomiting and weak pulse
- d. Both a and b
- e. All of the above

### 4. How do you determine if a person is having an anaphylactic reaction?

- a. Look for an allergic reaction
- b. Look for symptoms from 2 or more body systems
- c. Look for symptoms that persist for more than 5 minutes
- d. None of the above

### 5. What are the common triggers of anaphylaxis?

- a. Certain Foods, Latex, Insect Stings
- b. Cotton, Plastic, Exercise
- c. Medication, Exercise
- d. Both a and c

#### Indications

EpiPen® (epinephrine injection) 0.3 mg and EpiPen Jr® (epinephrine injection) 0.15 mg Auto-Injectors are for the emergency treatment of life-threatening allergic reactions (anaphylaxis) caused by allergens, exercise, or unknown triggers; and for people who are at increased risk for these reactions. EpiPen® and EpiPen Jr® are intended for immediate administration as emergency supportive therapy only. Seek immediate emergency medical help right away.

#### Important Safety Information

EpiPen® and EpiPen Jr® Auto-Injectors contain a single dose of epinephrine, which you (or your caregiver or others who may be in a position to administer EpiPen® or EpiPen Jr®) inject into the middle of your outer thigh (upper leg) (through clothing, if necessary). Get emergency medical help right away.

**Please see Important Safety Information at the end of this quiz. Please see accompanying full Prescribing Information and Patient Information.**



FEATURING

EPIPEN®

EPIPEN Jr®

(epinephrine injection) Auto-Injectors 0.3/0.15mg

# Mylan On Location™

## Anaphylaxis and Administering EpiPen® (epinephrine injection) Auto-Injector Quiz

6. Which of the following is a less common cause of anaphylaxis?
  - a. Pollen
  - b. Wasp sting
  - c. Aspirin
  - d. Latex gloves
  
7. When should you first administer epinephrine?
  - a. After other medications fail
  - b. Ten or more minutes after symptoms begin
  - c. At the first sign of symptoms
  - d. None of the above
  
8. What does the “R” mean in REACT?
  - a. Reach for epinephrine
  - b. Recognize signs and symptoms
  - c. Run for help
  - d. None of the above
  
9. After you “A”, administer Epinephrine, you should:
  - a. Leave the person alone
  - b. Call 911 or emergency responders
  - c. Stay with the person and treat again, if symptoms worsen
  - d. Both b and c
  
10. Where should epinephrine be administered?
  - a. In the outer thigh
  - b. In the arm
  - c. In the buttocks
  - d. In the lower leg
  
11. If epinephrine is administered and symptoms persist after 5 minutes:
  - a. Call 911
  - b. Monitor symptoms
  - c. Remove the allergen/trigger
  - d. Administer a second dose

### Important Safety Information (continued)

You may need further medical attention. Only a healthcare professional should give additional doses of epinephrine if you need more than two injections for a single anaphylactic episode. DO NOT INJECT INTO YOUR VEINS, BUTTOCKS, FINGERS, TOES, HANDS OR FEET. In case of accidental injection, please seek immediate medical treatment. Epinephrine should be used with caution if you have heart disease or are taking certain medicines that can cause heart-related (cardiac) symptoms.

**Please see Important Safety Information at the end of this quiz. Please see accompanying full Prescribing Information and Patient Information.**

# Mylan On Location™

## Anaphylaxis and Administering EpiPen® (epinephrine injection) Auto-Injector Quiz

12. A second reaction can occur within 1 to 72 hours.

- a. True
- b. False

13. EpiPen® should be stored:

- a. In the refrigerator
- b. In the carrier tube
- c. Away from light
- d. a and b
- e. b and c

14. EpiPen® Auto-Injector is approved for:

- a. Adults only
- b. Children 12+ years old and adults
- c. Anyone 66 lbs or more
- d. Anyone 50 lbs or more
- e. None of the above

15. EpiPen Jr® Auto-Injector is approved for:

- a. Children 5-11 years old
- b. Any child
- c. Anyone 25 lbs to 50 lbs
- d. Anyone 33 lbs to 66 lbs
- e. Anyone 25 lbs to 60 lbs

### Important Safety Information (continued)

Tell your doctor if you have certain medical conditions such as asthma, depression, thyroid disease, Parkinson's disease, diabetes, high blood pressure or heart disease, have any other medical conditions, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. Be sure to also tell your doctor all the medicines you take, especially medicines for asthma. **If you have certain medical conditions, or take certain medicines, your condition may get worse or you may have longer lasting side effects when you use EpiPen® or EpiPen Jr®.**

**Please see Important Safety Information at the end of this quiz. Please see accompanying full Prescribing Information and Patient Information.**

# Important Safety Information

## Important Safety Information

EpiPen® (epinephrine injection) 0.3 mg and EpiPen Jr® (epinephrine injection) 0.15 mg Auto-Injectors contain a single dose of epinephrine, which you (or your caregiver or others who may be in a position to administer EpiPen® or EpiPen Jr®) inject into the middle of your outer thigh (upper leg) (through clothing, if necessary). Get emergency medical help right away. You may need further medical attention. Only a healthcare professional should give additional doses of epinephrine if you need more than two injections for a single anaphylactic episode. **DO NOT INJECT INTO YOUR VEINS, BUTTOCKS, FINGERS, TOES, HANDS OR FEET.** In case of accidental injection, please seek immediate medical treatment. Epinephrine should be used with caution if you have heart disease or are taking certain medicines that can cause heart-related (cardiac) symptoms.

Tell your doctor if you have certain medical conditions such as asthma, depression, thyroid disease, Parkinson's disease, diabetes, high blood pressure or heart disease, have any other medical conditions, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. Be sure to also tell your doctor all the medicines you take, especially medicines for asthma. **If you have certain medical conditions, or take certain medicines, your condition may get worse or you may have longer lasting side effects when you use EpiPen® or EpiPen Jr®.**

The most common side effects may include increase in heart rate, stronger or irregular heartbeat, sweating, nausea or vomiting, difficulty breathing, paleness, dizziness, weakness, shakiness, headache, apprehension, nervousness or anxiety. These side effects may go away if you rest. **Tell your healthcare professional if you have any side effect that bothers you or that does not go away.**

**Please see the full Prescribing Information and Patient Information enclosed.**

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

For additional information, please contact us at 800-395-3376.



EpiPen® and EpiPen Jr® are registered trademarks owned by the Mylan companies. Mylan On Location™ is a trademark of Mylan Inc. The Mylan logo is a registered trademark of Mylan Inc.

© 2016 Mylan Specialty L.P. All rights reserved.  
9/15 EPI-2015-0819



FEATURING

**EPIPEN®** **EPIPEN Jr®**  
(epinephrine injection) Auto-Injectors 0.3/0.15mg

# Mylan On Location™

## Anaphylaxis and Administering EpiPen® (epinephrine injection) Auto-Injector Quiz

### Answer Key

1. c.
2. c.
3. e.
4. b.
5. d.
6. a.
7. c.
8. b.
9. d.
10. a.
11. d.
12. a.
13. e.
14. c.
15. d.