STATE OF TENNESSEE
DEPARTMENT OF CONSERVATION
DIVISION OF GEOLOGY

PREPARED IN COOPERATION WITH
UNITED STATES DEPARTMENT OF INTERIOR
GEOLOGICAL SURVEY

BULLETIN 58
PLATE 4, GREENEVILLE



EXPLANATION

- Mp — Pennington formation
- Mn — Newman limestone
- Mg — Grainger Formation
- MDs — Basal Mississippian and Devonian shale
- Sc — Clinch sandstone
- Oj — Juniata formation
- Omb — Martinsburg shale
- Olmc — Lower and middle parts of Chickamauga limestone
- Om — Moccasin formation
- Ob — Bays formation
- Och 2 — Chickamauga limestone, unit 2
- Och 1 — Chickamauga limestone, unit 1
- Osv — Sevier shale
- OCk — Knox dolomite or group, undivided
- Oma — Mascot dolomite
- Ok — Kingsport formation
- On — Newala formation
- Olv — Longview dolomite
- Oc — Chepultepec dolomite
- Ojb — Jonesboro limestone
- Ꞓcr — Copper Ridge dolomite
- Ꞓcc — Conococheague limestone
- Ꞓmn — Maynardville limestone
- Ꞓn — Nolichucky shale
- Ꞓcu — Upper Cambrian part of Conasauga group
- Ꞓrm — Rutledge, Rogersville, and Maryville formations, undivided
- Ꞓpv — Pumpkin Valley shale
- Ꞓcm — Middle Cambrian part of Conasauga group
- Ꞓhk — Honaker dolomite
- Ꞓr — Rome formation
- Ꞓs — Shady dolomite
- Ꞓe — Erwin formation
- Ꞓh — Hampton formation
- Ꞓu — Unicoi formation
- ocu — Ocoee series, undivided
- ocss — Sandsuck shale
- ocsb — Snowbird formation
- pꞒc — Pre-Cambrian crystalline complex
- Contact
- Fault

For master explanation see plate 11



LOCATION OF PLATES 1 TO 14

Base compiled and materials assembled by C. H. Heins, G. D. DeBuchananne and R. M. Richardson of the U. S. Geological Survey

Maps prepared for printing by N. B. Johnson, C. E. James, J. B. Canter, and others of the U. S. Geological Survey

Geology compiled by John Rodgers, U. S. Geological Survey
(For Description of sources see text)

GEOLOGIC MAP OF EAST TENNESSEE: GREENEVILLE

Scale $\frac{1}{125,000}$

1 0 5 10 Miles