MCH Services Title V Block Grant - Measurement Snapshot

Women and Maternal Health NPM 14.1 NVSS Percent of women with a past year preventive care visit NPM 14.1 NVSS Percent of women with a past year preventive care visit NPM 14.1 NVSS Percent of women with some with recompany of the preventive care visit NPM 14.1 NVSS Rate of severe maternal morbidity per 10,000 delivery hospitalizations NOM 10 PRAMS NOM 10 PRAMS Percent of infants born with feetal alcohol exposure in the last 3 months of pregnancy NOM 11 SID Rate of infants born with neonatal abstinence syndrome per 1,000 hospital births NOM 23 NVSS Rate of infants born to teen mothers per 1,000 females Percent of women who experience postpartum depressive symptoms following a recent live birth Percent of women who experience postpartum depressive symptoms following a recent live birth Percent of infants placed to sleep on their backs NPM 55 PRAMS Percent of infants placed to sleep on their backs NPM 55 PRAMS Percent of infants placed to sleep without sort objects or loose bedding Percent of infants placed to sleep without sort objects or loose bedding NPM 50 PRAMS Percent of low birth weight deliveries cle2500 grams) NOM 6 NVSS Percent of early term births (27, 38 weeks) NOM 8 NVSS Rate of perinatal mortality per 1,000 live births NOM 9.2 NVSS Rate of infant mortality per 1,000 live births NOM 9.3 NVSS Rate of neonatal mortality per 1,000 live births NOM 9.4 NVSS Rate of post neonatal mortality rate per 1,000 live births NOM 9.5 NVSS Rate of preterm-related wordality per 1,000 live births NOM 9.4 NVSS Rate of preterm-related mortality per 1,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 1,000 live births NOM 9.5 NVSS Rate of Precent of children without special health care needs having a medical home Percent of children who are physically active at least 60 minutes per day NPM 8.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 12 NSCH Percent of children who are physically active at least 60 minutes per day NOM	Measure	Data Source	Description	
NOM 1.1. NVSS. Percent of women who smoke during pregnancy				
NOM 3 NVSS Rate of severe maternal morbidity per 10,000 delivery hospitalizations NOM 10 PRAMS Percent of infants born with fetal alcohol exposure in the last 3 months of pregnancy NOM 11 SID Rate of infants born with fetal alcohol exposure in the last 3 months of pregnancy NOM 12 NVSS Rate of infants born with neonatal abstinence syndrome per 1,000 hospital births NOM 23 NVSS Rate of infants born to teen mothers per 1,000 females Percent of women who experience postpartum depressive symptoms following a recent live birth Perinatal and Infant Health NPM 59 PRAMS Percent of infants placed to sleep on their backs NPM 50 PRAMS Percent of infants placed on a separate sleep surface NPM 50 PRAMS Percent of infants placed to sleep on their backs NPM 50 PRAMS Percent of infants placed to sleep on their backs NPM 50 PRAMS Percent of infants placed to sleep without soft objects or loose bedding NPM 50 PRAMS Percent of infants placed to sleep without soft objects or loose bedding NPM 50 PRAMS Percent of fearly term births (437 weeks) NOM 6 NVSS Percent of perterm births (437 weeks) NOM 8 NVSS Percent of rearly term births (37, 38 weeks) NOM 8 NVSS Percent of rearly term births (37, 38 weeks) NOM 9.1 NVSS Rate of perinatal mortality per 1,000 live births NOM 9.2 NVSS Rate of infant mortality per 1,000 live births NOM 9.3 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.4 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.5 NVSS Rate of Percent of children receiving a developmental screening using a parent-completed screening tool NPM 1.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 1.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 1.1 NSCH Percent of children who live in households where someone smokes Percent of children without special health care needs having a medical home NPM 1.2 NSCH Percent of children who live in households where someone smokes NPM 1.1 NSCH Percent of children without	NPM 1	BRFSS	Percent of women with a past year preventive care visit	
NOM 10 PRAMS Percent of infants born with fetal alcohol exposure in the last 3 months of pregnancy NOM 11 SID Rate of infants born with the anala abstinence syndrome per 1,000 hospital births NOM 23 NVSS Rate of infants born to teen mothers per 1,000 females NOM 24 PRAMS Percent of women who experience postpartum depressive symptoms following a recent live birth Perinatal and Infant Health NPM 59 PRAMS Percent of infants placed to sleep on their backs NPM 50 PRAMS Percent of infants placed to sleep on their backs NPM 50 PRAMS Percent of infants placed to sleep without soft objects or loose bedding NPM 50 PRAMS Percent of infants placed to sleep without soft objects or loose bedding NPM 50 PRAMS Percent of for both birth weight deliveries (<2500 grams) NPM 50 PRAMS Percent of present births (<37 weeks) NPM 50 PRAMS Percent of present births (<37 weeks) NPM 50 Percent of present births (<37 weeks) NPM 51 NVSS Rate of present births (<37 weeks) NPM 52 Percent of present births (<37 weeks) NPM 53 Percent of present births (<37 weeks) NPM 54 NVSS Rate of present births (<37 weeks) NPM 55 Rate of present births (<37 weeks) NPM 50 Percent of contact mortality per 1,000 live births NPM 50 Percent of contact mortality per 1,000 live births NPM 50 Percent of contact mortality per 100,000 live births NPM 50 Percent of contact mortality per 100,000 live births NPM 51 NSS Rate of present births (solded unexplained infant Death (SUID) per 100,000 live births NPM 51 NSS Rate of Present of children without special beath care needs who received services necessary to make transitions to adult health care NPM 11 NSCH Percent of children who are physically active at least 60 minutes per day NPM 51 NSCH Percent of children who are very good health NPM 52 NSS Rate of hild mortality per 100,000 NPM 52 NSS Rate of	NPM 14.1	NVSS	Percent of women who smoke during pregnancy	
NOM 10 PRAMS Percent of infants born with feetal alcohol exposure in the last 3 months of pregnancy	NOM 2	SID	Rate of severe maternal morbidity per 10,000 delivery hospitalizations	
NOM 23 NVSS Rate of infants born with neonatal abstinence syndrome per 1,000 hospital births	NOM 3	NVSS	Rate of maternal mortality per 100,000 live births	
NOM 23 NVSS Rate of infants born to teen mothers per 1,000 females Pericant of women who experience postpartum depressive symptoms following a recent live birth Periatal and Infant Health NPM 5a PRAMS Percent of infants placed to sleep on their backs NPM 5b PRAMS Percent of infants placed to sleep on their backs NPM 5c PRAMS Percent of infants placed to sleep on their backs NPM 5c PRAMS Percent of infants placed to sleep without soft objects or loose bedding NOM 4 NVSS Percent of periate mortality per 1,000 live births NOM 5 NVSS Percent of periater mortality per 1,000 live births plus fetal deaths NOM 8 NVSS Percent of periater mortality per 1,000 live births NOM 9.1 NVSS Rate of perinatal mortality per 1,000 live births NOM 9.2 NVSS Rate of infant mortality per 1,000 live births NOM 9.3 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.4 NVSS Rate of periater mortality per 1,000 live births NOM 9.5 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.5 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.6 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.7 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.8 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.8 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.9 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.1 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.2 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.4 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.5 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.6 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.7 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.8 NVSS Rate of children who are wealth of the post live live live live live live live live	NOM 10	PRAMS	Percent of infants born with fetal alcohol exposure in the last 3 months of pregnancy	
PRAMS Percent of women who experience postpartum depressive symptoms following a recent live birth Perinatal and Infant Health NPM 5b PRAMS Percent of infants placed to sleep on their backs NPM 5b PRAMS Percent of infants placed to sleep on their backs NPM 5c PRAMS Percent of infants placed to sleep without soft objects or loose bedding NOM 4 NVS Percent of low birth weight deliveries (<2500 grams) NOM 5 NVS Percent of preterm births (<37 weeks) NOM 6 NVSS Percent of preterm births (<37 weeks) NOM 8 NVSS Percent of early term births (37, 38 weeks) NOM 9.1 NVSS Rate of perinatal mortality per 1,000 live births plus fetal deaths NOM 9.2 NVSS Rate of infant mortality per 1,000 live births NOM 9.3 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.4 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.5 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.4 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained Infant Death (SUID) per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained Infant Death (SUID) per 100,000 live births NPM 7.1 SID Rate of Sleep-related Sudden Unexplained Infant Death (SUID) per 100,000 live births NPM 8.1 NSCH Percent of children web are physically active at least 60 minutes per day NPM 11 NSCH Percent of children without special health care needs having a medical home Percent of children without special health care needs who received services necessary to make transitions to adult health care NOM 13 NSCH Percent of children with a metal/behavioral condition who receive treatment or counseling NOM 15 NSS Rate of Another with a metal/behavioral condition who receive treatment or counseling NOM 16 NSCH Percent of children with a metal/behavioral condition who receive treatment or counseling NOM 17 Percent of children with a revenue given to obese (BMI at or above 85th percentile) NOM 18 NSCH Percent of children with a revenue given to obese (BMI at o	NOM 11	SID	Rate of infants born with neonatal abstinence syndrome per 1,000 hospital births	
Perinatal and Infant Health Percent of infants placed to sleep on their backs NPM 5c PRAMS Percent of infants placed on a separate sleep surface NPM 5c PRAMS Percent of infants placed on a separate sleep surface NPM 5c PRAMS Percent of infants placed on a separate sleep surface NPM 5c PRAMS Percent of infants placed to sleep without soft objects or loose bedding NOM 4 NVSS Percent of low birth weight deliveries (<2500 grams) NOM 5 NVSS Percent of early term births (<37 weeks) NOM 6 NVSS Percent of early term births (37, 38 weeks) NOM 9.1 NVSS Rate of perinatal mortality per 1,000 live births plus fetal deaths NOM 9.1 NVSS Rate of infant mortality per 1,000 live births NOM 9.2 NVSS Rate of infant mortality per 1,000 live births NOM 9.3 NVSS Rate of post neonatal mortality reat per 1,000 live births NOM 9.3 NVSS Rate of preterm-related mortality per 1,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 1,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 1,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 1,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 1,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of hospitalization for non-fatal injury per 100,000 children NPM 9.6 NSCH Percent of children who are physically active at least 60 minutes per day NPM 11 NSCH Percent of children who are physically active at least 60 minutes per day NPM 12 NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14b NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 15 NSCH Percent of children who are physically active at least 60 minutes per day NOM 16 NSCH Percent of children with a revellent or very good health N	NOM 23	NVSS	Rate of infants born to teen mothers per 1,000 females	
NPM 5b PRAMS Percent of infants placed to sleep on their backs NPM 5b PRAMS Percent of infants placed to sleep on their backs NPM 5b PRAMS Percent of infants placed to sleep without soft objects or loose bedding NOM 4 NVSS Percent of preterm births (37 weeks) NOM 5 NVSS Percent of preterm births (37 weeks) NOM 6 NVSS Percent of preterm births (37 weeks) NOM 8 NVSS Percent of preterm births (37 weeks) NOM 9.1 NVSS Rate of perinatal mortality per 1,000 live births plus fetal deaths NOM 9.1 NVSS Rate of perinatal mortality per 1,000 live births plus fetal deaths NOM 9.2 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.3 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.4 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.5 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 1,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 1,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 10,000 live births NOM 9.5 NVSS Rate of preterm-related sudden Unexplained Infant Death (SUID) per 100,000 live births NOM 9.5 NVSS Rate of preterm-related sudden Unexplained Infant Death (SUID) per 100,000 live births NOM 9.5 NVSS Rate of preterm-related sudden Unexplained Infant Death (SUID) per 100,000 live births NOM 9.5 NVSS Rate of of Sleep-related Sudden Unexplained Infant Death (SUID) per 100,000 live births NOM 1.5 NSCH Percent of children without special health care needs having a medical home NPM 1.1 NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 1.2 NSCH Percent of children who live in households where someone smokes NOM 1.5 NSS Rate of children meeting the criteria developed for school readiness NOM 1.5 NSS Rate of children with a mental/behavioral condition who received services necessary to make transitions to adult health care need shaving a medical home NPM 1.2 NSCH Percent of children with a percent of	NOM 24	PRAMS	Percent of women who experience postpartum depressive symptoms following a recent live birth	
NPM 5b PRAMS Percent of infants placed to seep without soft objects or loose bedding NPM 5c PRAMS Percent of for birth weight deliveries (<2500 grams) NOM 4 NVSS Percent of for birth weight deliveries (<2500 grams) NOM 5 NVSS Percent of for birth weight deliveries (<2500 grams) NOM 6 NVSS Percent of early term births (37 weeks) NOM 6 NVSS Percent of early term births (37, 38 weeks) NOM 9.0 NVSS Rate of prefeatal mortality per 1,000 live births plus fetal deaths NOM 9.1 NVSS Rate of infant mortality per 1,000 live births NOM 9.2 NVSS Rate of infant mortality per 1,000 live births NOM 9.3 NVSS Rate of prefeatal mortality per 1,000 live births NOM 9.3 NVSS Rate of prefeatal mortality per 1,000 live births NOM 9.5 NVSS Rate of preferm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preferm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preferm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preferm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preferm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preferm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preferm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preferm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preferm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preferm related with self-preferming using a parent-completed screening tool NPM 7.1 SID Reterent of children who are physically active at least 60 minutes per day NPM 1.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 1.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 1.2 NSCH Percent of children who are physically active at least 60 minutes per day NPM 1.2 NSCH Percent of children who are physically active at least 60 minutes per day NPM 1.2 NSCH Percent of children who are physically active at least 60 minutes per day NPM 1.2 NSCH Percent of children who are prevented benefits per 100,000 NPM				
NPM 5C PRAMS Percent of infants placed to sleep without soft objects or loose bedding NOM 4 NVSS Percent of low birth weight deliveries (<2500 grams) NOM 5 NVSS Percent of John birth weight deliveries (<2500 grams) NOM 6 NVSS Percent of percent of joretrom births (<37 weeks) NOM 8 NVSS Rate of perinatal mortality per 1,000 live births plus fetal deaths NOM 9.1 NVSS Rate of infant mortality per 1,000 live births NOM 9.2 NVSS Rate of neonatal mortality per 1,000 live births NOM 9.3 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.3 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.4 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained Infant Death (SUID) per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained Infant Death (SUID) per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained Infant Death (SUID) per 100,000 live births NPM 1.1 NSCH Percent of children receiving a developmental screening using a parent-completed screening tool NPM 1.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 1.1 NSCH Percent of children without special health care needs having a medical home NPM 1.2 NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 1.4 NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 1.3 NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 1.5 NSCH Percent of children with a mental/behavioral condition who received treatment or counseling NOM 1.5 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 1.5 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 1.5 NSCH Percent of children w	NPM 5a	PRAMS	Percent of infants placed to sleep on their backs	
NOM 4 NVSS Percent of low birth weight deliveries (<2500 grams) NOM 5 NVSS Percent of eriths (<37 weeks) NOM 6 NVSS Percent of preterm births (<37 weeks) NOM 8 NVSS Percent of eriths (<37 weeks) NOM 9.1 NVSS Rate of perinatal mortality per 1,000 live births plus fetal deaths NOM 9.1 NVSS Rate of infant mortality per 1,000 live births NOM 9.2 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.3 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.4 NVSS Rate of post neonatal mortality per 10,000 live births NOM 9.5 NVSS Rate of Seper-related Sudden Unexplained lifant Death (SUID) per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained lifant Death (SUID) per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained lifant Death (SUID) per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained lifant Death (SUID) per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained lifant Death (SUID) per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained lifant Death (SUID) per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained lifant Death (SUID) per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained lifant Death (SUID) per 100,000 live births NOM 1.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 1.1 NSCH Percent of children who are physically active at least 60 minutes per day NOM 1.5 NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NOM 1.5 NSCH Percent of children meeting the criteria developed for school readiness NOM 1.5 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 1.5 NSCH Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 1.6 NSCH Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 1.6	NPM 5b	PRAMS	Percent of infants placed on a separate sleep surface	
NOM 5 NVSS Percent of preterm births (<37 weeks) NOM 6 NVSS Percent of early term births (37, 38 weeks) NOM 8 NVSS Rate of perinatal mortality per 1,000 live births plus fetal deaths NOM 9.1 NVSS Rate of infant mortality per 1,000 live births NOM 9.2 NVSS Rate of neonatal mortality per 1,000 live births NOM 9.3 NVSS Rate of neonatal mortality per 1,000 live births NOM 9.4 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.5 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.6 NVSS Rate of post neonatal mortality per 100,000 live births NOM 9.7 NVSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained Infant Death (SUID) per 100,000 live births NPM 8.1 NVSCH Percent of children receiving a developmental screening using a parent-completed screening tool NPM 8.1 NVSCH Percent of children without special health care needs having a medical home NPM 11 NVSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14 NVSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14 NVSCH Percent of children who live in households where someone smokes NOM 13 NVSCH Percent of children meeting the criteria developed for school readiness NOM 15 NVSS Rate of child mortality per 100,000 NOM 18 NVSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NVSCH Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 20 NVSCH Rate of hospitalization for non-fatal injury per 100,000 NOM 16.2 NVSCH Rate of hospitalization for non-fatal injury per 100,000 NOM 16.2 NVSS Rate of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortal with special health care needs	NPM 5c	PRAMS	Percent of infants placed to sleep without soft objects or loose bedding	
NOM 6 NVSS Percent of early term births (37, 38 weeks) NOM 8 NVSS Rate of perinatal mortality per 1,000 live births plus fetal deaths NOM 9.1 NVSS Rate of infant mortality per 1,000 live births NOM 9.2 NVSS Rate of nonatal mortality per 1,000 live births NOM 9.3 NVSS Rate of post neonatal mortality per 10,000 live births NOM 9.4 NVSS Rate of post neonatal mortality per 10,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 10,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained Infant Death (SUID) per 100,000 live births Child Health NPM 6 NSCH Percent of children receiving a developmental screening using a parent-completed screening tool NPM 7.1 SID Rate of hospitalization for non-fatal injury per 100,000 children NPM 8.1 NSCH Percent of children without special health care needs having a medical home Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14b NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14b NSCH Percent of children without with in households where someone smokes NOM 13 NSCH Percent of children with per 100,000 NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 20 WIC Percent of children in excellent or very good health NOM 20 NSCH Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 16.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NOM 16.3 NSS Rate of adolescents who are physically active at least 60 minutes per day NOM 16.3 NVSS Rate of adolescent mortality per 100,000 NOM 16.3 NVSS Rate of adolescent mortality per 100,000 NOM 16.4 NSCH Percent of children with special health care needs having a m	NOM 4	NVSS	Percent of low birth weight deliveries (<2500 grams)	
NOM 8 NVSS Rate of perinatal mortality per 1,000 live births plus fetal deaths NOM 9.1 NVSS Rate of infant mortality per 1,000 live births NOM 9.2 NVSS Rate of neonatal mortality per 1,000 live births NOM 9.3 NVSS Rate of perinatal mortality per 1,000 live births NOM 9.4 NVSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 100,000 live births **Child Health** NPM 6 NSCH Percent of children receiving a developmental screening using a parent-completed screening tool NPM 7.1 SID Rate of hospitalization for non-fatal injury per 100,000 children NPM 8.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 11 NSCH Percent of children without special health care needs having a medical home Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14b NSCH Percent of children with olive in households where someone smokes NOM 13 NSCH Percent of children meeting the criteria developed for school readiness NOM 15 NSCS Rate of child mortality per 100,000 NOM 18 NSCH Percent of children in excellent or very good health NOM 20 WIC Percent of children in excellent or very good health NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Rate of hospitalization for non-fatal injury per 100,000 NOM 16.2 NVSS Rate of adolescent swind are physically active at least 60 minutes per day NOM 16.3 NVSS Rate of adolescent swind are physically active at least 60 minutes per day NOM 16.3 NVSS Rate of adolescent swind are physically active at least 60 minutes per day NOM 16.3 NVSS Rate of adolescent swind are physically active at least 60 minutes per day NOM 16.4 NVSS Rate of adolescent swind are physically active at least 60 minutes per day NOM 16.5 NVSS Rate of	NOM 5	NVSS	Percent of preterm births (<37 weeks)	
NOM 8 NVSS Rate of perinatal mortality per 1,000 live births plus fetal deaths NOM 9.1 NVSS Rate of infant mortality per 1,000 live births NOM 9.2 NVSS Rate of neonatal mortality per 1,000 live births NOM 9.3 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.4 NVSS Rate of preterm-related mortality per 1,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of preterm-related mortality per 100,000 children NPM 9.1 NSCH Percent of children receiving a developmental screening using a parent-completed screening tool NPM 7.1 SID Rate of hospitalization for non-fatal injury per 100,000 children NPM 1.1 NSCH Percent of children without special health care needs having a medical home Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14b NSCH Percent of children who live in households where someone smokes NOM 13 NSCH Percent of children meeting the criteria developed for school readiness NOM 15 NVSS Rate of child mortality per 100,000 NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children who are overweight or obese (BMI at or above 85th percentile) Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Rate of hospitalization for non-fatal injury per 100,000 NOM 16.2 NVSS Rate of adolescent swind are physically active at least 60 minutes per day NOM 16.3 NVSS Rate of adolescent swind are physically active at least 60 minutes per day NOM 16.3 NVSS Rate of adolescent swind are physically active at least 60 minutes per day NOM 16.3 NVSS Rate of adolescent s	NOM 6	NVSS	Percent of early term births (37, 38 weeks)	
NOM 9.1 NVSS Rate of infant mortality per 1,000 live births NOM 9.2 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.3 NVSS Rate of post neonatal mortality per 1,000 live births NOM 9.4 NVSS Rate of post neonatal mortality per 10,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained Infant Death (SUID) per 100,000 live births NPM 6 NSCH Percent of children receiving a developmental screening using a parent-completed screening tool NPM 7.1 SID Rate of hospitalization for non-fatal injury per 100,000 children NPM 8.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 11 NSCH Percent of children without special health care needs having a medical home Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14b NSCH Percent of children who live in households where someone smokes NOM 13 NSCH Percent of children who live in households where someone smokes NOM 18 NSCH Percent of children meeting the criteria developed for school readiness NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 20 NSCH Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 21 NSCH Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 22 NSCH Rate of hospitalization for non-fatal injury per 100,000 NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent mortality per 100,000 NOM 16.3 NVSS Rate of adolescent mort vehicle mortality per 100,000 NOM 16.4 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 20 YRBSS Percent of children with special health care needs having a medical home Percent of children with special health care needs having a medical home	NOM 8	NVSS		
NOM 9.2 NVSS Rate of neonatal mortality per 1,000 live births NOM 9.3 NVSS Rate of post neonatal mortality rate per 1,000 live births NOM 9.4 NVSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained Infant Death (SUID) per 100,000 live births Child Health NPM 6 NSCH Percent of children receiving a developmental screening using a parent-completed screening tool NPM 7.1 SID Rate of hospitalization for non-fatal injury per 100,000 children NPM 8.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 11 NSCH Percent of children without special health care needs having a medical home NPM 12 NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14b NSCH Percent of children who live in households where someone smokes NOM 13 NSCH Percent of children meeting the criteria developed for school readiness NOM 15 NNSS Rate of child mortality per 100,000 NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Rate of hospitalization for non-fatal injury per 100,000 NOM 16.1 NVSS Rate of adolescents who are physically active at least 60 minutes per day NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 20 YRBSS Percent of children with special health care needs having a medical home Percent of children with special health care needs having a medical home Percent of children with special health care needs baving a redical home Percent of children with special health care needs who received services necessary to make transitions to adult health care	NOM 9.1	NVSS		
NOM 9.3 NVSS Rate of post neonatal mortality rate per 1,000 live births NOM 9.4 NVSS Rate of Steepr-related mortality per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained Infant Death (SUID) per 100,000 live births Child Health NPM 6 NSCH Percent of children receiving a developmental screening using a parent-completed screening tool NPM 7.1 SID Rate of hospitalization for non-fatal injury per 100,000 children NPM 8.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 8.1 NSCH Percent of children without special health care needs having a medical home NPM 12 NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14b NSCH Percent of children meeting the criteria developed for school readiness NOM 13 NSCH Percent of children meeting the criteria developed for school readiness NOM 15 NVSS Rate of child mortality per 100,000 NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 26 Percent of children who are not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent mortality per 100,000 NOM 16.3 NVSS Rate of adolescent mortality per 100,000 NOM 16.4 Percent of children with special health care needs having a medical home Children and Youth with Special Health Care Needs Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs who received services necessary to make transitions to adult health care	NOM 9.2			
NOM 9.4 NOSS Rate of preterm-related mortality per 100,000 live births NOM 9.5 NVSS Rate of Sleep-related Sudden Unexplained Infant Death (SUID) per 100,000 live births Child Health NPM 6 NSCH Percent of children receiving a developmental screening using a parent-completed screening tool NPM 7.1 SID Rate of hospitalization for non-fatal injury per 100,000 children NPM 8.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 11 NSCH Percent of children without special health care needs having a medical home Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14b NSCH Percent of children who live in households where someone smokes NOM 13 NSCH Percent of children meeting the criteria developed for school readiness NOM 15 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children in excellent or very good health NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who were not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NOM 16.1 NVSS Rate of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent mortality per 100,000 NOM 16.3 NVSS Rate of adolescent mortality per 100,000 NOM 20 YRBSS Percent of children with special health care needs having a medical home Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs who received services necessary to make transitions to adult health care Percent of children with special health care needs who received services necessary to make transitions to adult health care				
NOM 9.5 NYSS Rate of Sleep-related Sudden Unexplained Infant Death (SUID) per 100,000 live births Child Health NPM 6 NSCH Percent of children receiving a developmental screening using a parent-completed screening tool NPM 7.1 SID Rate of hospitalization for non-fatal injury per 100,000 children NPM 8.1 NSCH Percent of children without special health care needs having a medical home Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 12 NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14b NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 15b NSCH Percent of children who live in households where someone smokes NOM 15 NVSS Rate of children meeting the criteria developed for school readiness NOM 15 NVSS Rate of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who were not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NPM 8.2 VRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 20 VRBSS Percent of children with special health care needs having a medical home Children and Youth with Special Health Care Needs Children and Youth with Special health care needs who received services necessary to make transitions to adult health care Adult health care NPM 12 NSCH Percent of children with special health ca	NOM 9.4	NVSS		
NPM 6 NSCH Percent of children receiving a developmental screening using a parent-completed screening tool NPM 7.1 SID Rate of hospitalization for non-fatal injury per 100,000 children NPM 8.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 1.1 NSCH Percent of children without special health care needs having a medical home NPM 12 NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14b NSCH Percent of children who live in households where someone smokes NOM 13 NSCH Percent of children meeting the criteria developed for school readiness NOM 15 NVSS Rate of child mortality per 100,000 NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children in excellent or very good health NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who were not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NPM 8.2 YRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 16.4 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NPM 12 NSCH Percent of children with special health care needs who received services necessary to make transitions to adult health care	NOM 9.5	NVSS	· · · · · · · · · · · · · · · · · · ·	
NPM 7.1 SID Rate of hospitalization for non-fatal injury per 100,000 children NPM 8.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 11 NSCH Percent of children without special health care needs having a medical home Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 12 NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14b NSCH Percent of children who live in households where someone smokes NOM 13 NSCH Percent of children meeting the criteria developed for school readiness NOM 15 NYSS Rate of child mortality per 100,000 NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who were not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NOM 16.1 NYSS Rate of adolescents who are physically active at least 60 minutes per day NOM 16.1 NYSS Rate of adolescent mortality per 100,000 NOM 16.2 NYSS Rate of adolescent mortality per 100,000 NOM 16.3 NYSS Rate of adolescent mortor vehicle mortality per 100,000 NOM 20 YRBSS Percent of children who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care				
NPM 7.1 SID Rate of hospitalization for non-fatal injury per 100,000 children NPM 8.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 11 NSCH Percent of children without special health care needs having a medical home Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 12 NSCH Percent of children who live in households where someone smokes NOM 13 NSCH Percent of children meeting the criteria developed for school readiness NOM 15 NVSS Rate of child mortality per 100,000 NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who were not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NOM 16.1 NVSS Rate of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent mortor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care	NPM 6	NSCH	Percent of children receiving a developmental screening using a parent-completed screening tool	
NPM 8.1 NSCH Percent of children who are physically active at least 60 minutes per day NPM 11 NSCH Percent of children without special health care needs having a medical home Percent of children without special health care needs who received services necessary to make transitions to adult health care Percent of children without special health care needs who received services necessary to make transitions to adult health care Percent of children who live in households where someone smokes NOM 13 NSCH Percent of children meeting the criteria developed for school readiness NOM 15 NVSS Rate of child mortality per 100,000 NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 16.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NOM 16.1 NVSS Rate of adolescents who are physically active at least 60 minutes per day NOM 16.2 NVSS Rate of adolescent mortality per 100,000 NOM 16.3 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 20 YRBSS Percent of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care Percent of children with special health care needs receiving care in a well-functioning system	NPM 7.1	SID		
NPM 11 NSCH Percent of children without special health care needs having a medical home NPM 12 NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14b NSCH Percent of children who live in households where someone smokes NOM 13 NSCH Percent of children meeting the criteria developed for school readiness NOM 15 NVSS Rate of child mortality per 100,000 NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children in excellent or very good health NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who were not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NPM 8.2 YRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent mortality per 100,000 NOM 16.3 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NPM 12 NSCH Percent of children with special health care needs who received services necessary to make transitions to adult health care	NPM 8.1	NSCH		
NPM 12 NSCH Percent of children without special health care needs who received services necessary to make transitions to adult health care NPM 14b NSCH Percent of children who live in households where someone smokes NOM 13 NSCH Percent of children meeting the criteria developed for school readiness NOM 15 NVSS Rate of child mortality per 100,000 NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 27 NSCH Rate of hospitalization for non-fatal injury per 100,000 NPM 8.2 YRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NPM 12 NSCH Percent of children with special health care needs who received services necessary to make transitions to adult health care	NPM 11	NSCH		
Adult health care NPM 14b NSCH Percent of children who live in households where someone smokes NOM 13 NSCH Percent of children meeting the criteria developed for school readiness NOM 15 NVSS Rate of child mortality per 100,000 NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who were not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NPM 8.2 YRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NPM 12 NSCH Percent of children with special health care needs receiving care in a well-functioning system				
NOM 13 NSCH Percent of children meeting the criteria developed for school readiness NOM 15 NVSS Rate of child mortality per 100,000 NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children in excellent or very good health NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who were not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NPM 8.2 YRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NPM 12 NSCH Percent of children with special health care needs receiving care in a well-functioning system	NPM 12	NSCH		
NOM 13 NSCH Percent of children meeting the criteria developed for school readiness NOM 15 NVSS Rate of child mortality per 100,000 NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children in excellent or very good health NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who were not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NPM 8.2 YRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NPM 12 NSCH Percent of children with special health care needs receiving care in a well-functioning system	NPM 14b	NSCH	Percent of children who live in households where someone smokes	
NOM 15 NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who were not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NPM 8.2 YRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NSCH Percent of children with special health care needs receiving care in a well-functioning system	NOM 13		Percent of children meeting the criteria developed for school readiness	
NOM 18 NSCH Percent of children with a mental/behavioral condition who receive treatment or counseling NOM 19 NSCH Percent of children in excellent or very good health NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who were not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NPM 8.2 YRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system	NOM 15	NVSS		
NOM 19 NSCH Percent of children in excellent or very good health NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who were not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NPM 8.2 YRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system				
NOM 20 WIC Percent of children who are overweight or obese (BMI at or above 85th percentile) NOM 25 NSCH Percent of children who were not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NPM 8.2 YRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system		NSCH		
NOM 25 NSCH Percent of children who were not able to obtain need health care in the last year Adolescent Health NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NPM 8.2 YRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system				
NPM 7.2 NSCH Rate of hospitalization for non-fatal injury per 100,000 NPM 8.2 YRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system				
NPM 8.2 YRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system			,	
NPM 8.2 YRBSS Percent of adolescents who are physically active at least 60 minutes per day NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system	NPM 7.2	NSCH	Rate of hospitalization for non-fatal injury per 100,000	
NOM 16.1 NVSS Rate of adolescent mortality per 100,000 NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system				
NOM 16.2 NVSS Rate of adolescent motor vehicle mortality per 100,000 NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home NPM 12 NSCH Percent of children with special health care needs who received services necessary to make transitions to adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system			, , , , ,	
NOM 16.3 NVSS Rate of adolescent suicide per 100,000 NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home NPM 12 NSCH Percent of children with special health care needs who received services necessary to make transitions to adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system				
NOM 20 YRBSS Percent of adolescents who are obese Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home Percent of children with special health care needs who received services necessary to make transitions to adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system				
Children and Youth with Special Health Care Needs NPM 11 NSCH Percent of children with special health care needs having a medical home NPM 12 NSCH Percent of children with special health care needs who received services necessary to make transitions to adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system				
NPM 11 NSCH Percent of children with special health care needs having a medical home NPM 12 NSCH Percent of children with special health care needs who received services necessary to make transitions to adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system				
NPM 12 NSCH Percent of children with special health care needs who received services necessary to make transitions to adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system	NPM 11	NSCH		
NOM 17.2 NSCH adult health care NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system				
NOM 17.2 NSCH Percent of children with special health care needs receiving care in a well-functioning system	NPM 12	NSCH		
	NOM 17.2	NSCH		
propries process of children who were not able to obtain need health care in the last year				
Legend	10141 23	1.13611		

Note: Color rating is based on statistically significant changes in the most recent data available

Rating: green=improving; yellow=no change; red=worsening; no color=insufficient data to establish trend

Data Source Definitions: BRFSS=Behavioral Risk Factor Surveillance System (2009-2017); NSCH=National Survey of Children's Health (2016-2017); NVSS=National Vital Statistics System (2009-2017); PRAMS=Pregnancy Risk Assessment Monitoring System (2008-2015); SID=State Inpatient Database (2008-2016); WIC=Special Supplemental Nutrition Program for Women, Infants, and Children (even years 2008-2014); YRBSS=Youth Risk Behavior Surveillance System (odd years 2005-2017)