

**** NOTE: Hearing results should be submitted on the Newborn Screening Filter Card OR carbon form
Use this form ONLY if these are not available & fax to Newborn Screening Program at 615-532-8555****

Tennessee Department of Health
Newborn Screening Follow Up Program
1st Floor, R.S. Gass Building
630 Hart Lane, Nashville, Tennessee 37243
Phone (855) 202-1357 Fax (615) 532-8555

Hearing Screen Only Form

(See note above – Filter Card or Carbon Copy Form should be used first if possible)

Child's Last Name First Name Middle Name Gender (Twin: A or B) Date of Birth

Birth Mother's Last Name First Name Maiden Name State Lab TDH#

Address City State/Zip Phone

Primary Care Provider Phone

Birth Hospital Name: _____ City/State: _____

If this infant was TRANSFERRED, list hospital: _____

Person filling out form (print name): _____ Phone: _____

Facility/Provider Name: _____ City: _____

RESULTS – INITIAL SCREEN:

Date of Initial Hearing Screen: ____/____/____

Method: ABR/AABR OAE

Results: R: Pass Refer L: Pass Refer

Risk Factors: Mark in box at bottom

IF INFANT DID NOT PASS INITIAL SCREEN AND FURTHER EVALUATION (RESCREEN OR DIAGNOSTIC) WAS DONE:

Test done by: Hospital PCP Audiologist ENT/Otolary Other _____

Date of Evaluation _____ Type of Evaluation: ABR/AABR OAE Tymp/Reflex ASSR Behavioral

Results: R: Pass Refer L: Pass Refer

IF THIS INFANT WAS REFERRED TO: Audiologist ENT Other _____

Name: _____ Phone: _____ Appointment on: ____/____/____

Comments/Reason: _____

Risk Factors: (see below, check all that apply)

1 2 3 4 5 6 7 A C D F

- 1. NICU >5 Days
- 2. Syndrome associated with progressive or late onset HL
- 3. Family History of permanent childhood hearing loss
- 4. Birth conditions or findings including microtia/atresia, ear dysplasia, cleft lip and/or palate, temporal bone abnormalities, white forelock, microphthalmia, congenital microcephaly, congenital or acquired hydrocephalus.
- 5. In-utero infections, such as CMV, Herpes, Rubella, Syphilis and Toxoplasmosis; Zika + Infant
- 6. ECMO
- 7. Asphyxia or Hypoxic Encephalopathy

- A. Events associated with hearing loss including significant head trauma, (especially basal skull/temporal bone fractures) or chemotherapy
- C. Aminoglycoside administration >5 days
- D. Hyperbilirubinemia requiring exchange transfusion
- F. Postnatal culture-positive infections associated with Sensorineural Hearing Loss, including confirmed bacterial and viral (especially Herpes virus and Varicella) meningitis and encephalitis.

