

**2024 PCMH
Quality Metrics and Thresholds – Family Practice**

Core metric	Description	Threshold
1. Antidepressant Medication Management (adults only) – Effective Continuation Phase	The percentage of members 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression and who remained on an antidepressant medication for at least 180 days (6 months).	≥ 40.00%
2. Breast Cancer Screening - Electronic	The percentage of members 50-74 years of age who were recommended for routine breast cancer screening and had a mammogram to screen for breast cancer.	≥ 47.00%
3. Controlling High Blood Pressure	The percentage of members 18–85 years of age who had a diagnosis of hypertension (HTN) and whose blood pressure (BP) was adequately controlled (<140/90 mm Hg) during the measurement year.	≥ 60.00%
4. Childhood Immunizations Status - Combination 10	The percentage of children 2 years of age who had four diphtheria, tetanus and acellular pertussis (DTaP); three polio (IPV); one measles, mumps and rubella (MMR); three haemophilus influenza type B (HiB); three hepatitis B (HepB), one chicken pox (VZV); four pneumococcal conjugate (PCV); one hepatitis A (HepA); two or three rotavirus (RV); and two influenza (flu) vaccines by their second birthday.	≥ 42.00%
5. Blood Pressure Control for Patients with Diabetes	The percentage of members 18–75 years of age with diabetes (types 1 and 2) whose blood pressure (BP) was adequately controlled (<140/90 mm Hg).	≥ 62.00%
6. Eye Exam for Patients with Diabetes	The percentage of members 18–75 years of age with diabetes (types 1 and 2) who had a retinal eye exam.	≥ 51.00%
7. Glycemic Status Assessment for Patients with Diabetes <ul style="list-style-type: none"> Glycemic Status <8.0% 	The percentage of members 18–75 years of age with diabetes (types 1 and 2) most recent glycemic status (hemoglobin A1c [HbA1c] or glucose management indicator [GMI]) was at the following levels during the measurement year: <ul style="list-style-type: none"> Glycemic Status <8.0%. 	≥ 47.00%
8. Child and Adolescent Well-Care Visits		
<ul style="list-style-type: none"> Ages 3 – 11 Years 	The percentage of members 3–11 years of age who had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner during the measurement year.	≥ 65.00%
<ul style="list-style-type: none"> Ages 12 -17 Years 	The percentage of members 12–17 years of age who had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner during the measurement year.	≥ 57.00%
<ul style="list-style-type: none"> Ages 18 – 21 Years 	The percentage of members 18–21 years of age who had at least one comprehensive well-care visit with a PCP or OB/GYN practitioner during the measurement year.	≥ 39.00%

Core metric	Description	Threshold
9. Well-Child Visits in the First 30 Months of Life		
<ul style="list-style-type: none"> Well-Child Visits in the First 15 Months 	<p>The percentage of members who had the following number of well-child visits with a PCP during the last 15 months.</p> <ul style="list-style-type: none"> Children who turned 15 months old during the measurement year: Six or more well-child visits. 	≥ 61.00%
<ul style="list-style-type: none"> Well-Child Visits for Age 15 Months - 30 Months 	<p>The percentage of members who had the following number of well-child visits with a PCP during the last 15 months.</p> <ul style="list-style-type: none"> Children who turned 30 months old during the measurement year: Two or more well-child visits. 	≥ 71.00%
10. Immunizations for Adolescents, Total Rate – Combination 2	<p>The percentage of adolescents 13 years of age who had one dose of meningococcal vaccine, one tetanus, diphtheria toxoids and acellular pertussis (Tdap) vaccine, and have completed the human papillomavirus (HPV) vaccine series by their 13th birthday.</p>	≥ 26.00%