

**2020 & 2021 Patient Centered Medical Home
Quality Metrics for Family Practices**

Core metric	Description	Threshold
1. Antidepressant Medication Management (adults only) – Effective Continuation Phase	The percentage of members 18 and older who were treated with antidepressant medication, had a diagnosis of major depression, and who remained on an antidepressant medication for at least 180 days (6 months).	≥ 40.00%
2. Asthma Medication Ratio	The percentage of members 5–64 years of age who were identified as having persistent asthma and had a ratio of controller medications to total asthma medications of 0.50 or greater during the measurement year.	≥ 81.00%
3. Controlling High Blood Pressure	The percentage of members 18–85 years of age who had a diagnosis of hypertension (HTN) and whose BP was adequately controlled (<140/90 mm Hg) during the measurement year.	≥ 49.00%
4. Childhood Immunizations Status – Combination 10	The percentage children 2 years of age who had 4 DTaP, 3 polio, 1 MMR, 3 HiB, 3 HepB, 1 VZV, 4 PCV, 1 HepA, 2 or 3 RV, and 2 influenza vaccines by their second birthday.	≥ 42.00%
5. Comprehensive Diabetes Care: BP control (<140/90 mm Hg)	The percentage of members 18- 75 years of age with diabetes (type 1 and type 2), whose most recent blood pressure reading is less than 140/90 mm Hg (controlled).	≥ 56.00%
6. Comprehensive Diabetes Care: Eye exam (retinal) performed	The percentage of members 18- 75 years of age with type diabetes (type 1 and type 2), who had an eye exam (retinal) performed.	≥ 51.00%
7. Comprehensive Diabetes Care: HbA1c poor control (>9.0%)	The percentage of members 18- 75 years of age with diabetes (type 1 and type 2), whose most recent HbA1c level during the measurement year is greater than 9.0%.	≤ 47.00%
8. Child and Adolescent Well-Care Visits		
• Ages 3 – 11 years	The percentage of members 3-11 years of age who had at least one comprehensive well-care visit with a PCP or OB/GYN practitioner during the measurement year.	≥ 65.00%
• Ages 12 – 17 years	The percentage of members 12-17 years of age who had at least one comprehensive well-care visit with a PCP or OB/GYN practitioner during the measurement year.	≥ 57.00%
• Ages 18 – 21 years	The percentage of members 18-21 years of age who had at least one comprehensive well-care visit with a PCP or OB/GYN practitioner during the measurement year.	≥ 39.00%
9. Well-Child Visits in the First 30 Months of Life		
• Well-Child Visits in the First 15 Months	The percentage of members who had the following number of well-child visits with a PCP during the last 15 months. Children who turned 15 months old during the measurement year: Six or more well-child visits.	≥ 61.00%
• Well-Child Visits for Age 15 Months – 30 Months	The percentage of members who had the following number of well-child visits with a PCP during the last 15 months. Children who turned 30 months old during the measurement year: Two or more well-child visits.	≥ 71.00%
10. Immunizations for Adolescents – Total rate-Combination 2	The percentage of adolescents 13 years of age who had one dose of meningococcal vaccine, one Tdap and have completed the human papillomavirus (HPV) vaccine series by their 13th birthday.	≥ 26.00%